

Amendments to the Claims:

1. (Currently Amended) A pair of primers for specifically amplifying an hsp 65 (Heat Shock Protein 65) gene fragment of mycobacterial species ~~comprising, wherein one primer comprises~~ the nucleotide ~~sequence sequences as shown in~~ of SEQ ID NO: 55 and ~~the other primer comprises the nucleotide sequence of~~ SEQ ID NO: 56.

2. (Currently Amended) A polynucleotide of an hsp 65 gene fragment of mycobacterial species, wherein the fragment is amplified by using a pair of primers specifically amplifying the hsp 65 gene fragment of mycobacterial species, ~~wherein comprising one primer comprises~~ the nucleotide ~~sequence sequences as shown in~~ of SEQ ID NO: 55 and ~~the other primer comprises the nucleotide sequence of~~ SEQ ID NO: 56.

3. (Currently Amended) A polynucleotide selected from the group of ~~polynucleotides consisting of polynucleotides as shown in~~ SEQ ID NO: 1 to SEQ ID NO: 54, and polynucleotides complementary thereto.

4. (Currently Amended) A polynucleotide set for the detection or identification of mycobacterial species wherein the set comprises at least an hsp 65 gene fragment selected from the group of ~~polynucleotides consisting of the polynucleotides as shown in~~ SEQ ID NO: 1 to SEQ ID NO: 54 and polynucleotides complementary thereto.

5. (Currently Amended) A method for the identification of mycobacterial species comprising the steps of:

(1) amplifying an hsp 65 gene fragment of mycobacterial species of interest with primers for specifically amplifying the hsp 65 gene fragment, ~~wherein one primer comprises the nucleotide sequence of~~ SEQ ID NO: 55 and ~~the other primer comprises the nucleotide sequence of~~ SEQ ID NO: 56;

(2) analyzing a nucleotide sequence of the amplified hsp 65 gene fragment; and

(3) comparing multi-aligning the nucleotide sequence of the amplified hsp 65 gene fragment obtained analyzed in step (2) with a 604-bp hsp 65 gene fragment of a reference mycobacterial species a polynucleotide set comprising at least one hsp 65 gene fragment selected from the group of polynucleotides consisting of SEQ ID NO: 1 to SEQ ID NO: 54 and polynucleotides complementary thereto, to infer a phylogenetic tree.

6. (Cancelled)

7. (Cancelled)

8. (Original) A method for the identification of mycobacterial species comprising the steps of:

(1) amplifying an hsp 65 gene fragment of mycobacterial species with primers of claim 1; and

(2) analyzing the amplified fragment according to the RFLP (Restriction Fragment Length Polymorphism) analysis method using a restriction enzyme recognition site in the amplified fragment.

9. (Original) The method of claim 8, wherein the restriction enzyme is Xho I.

10. (Currently Amended) The method of claim 9 comprising the step of treating the amplified hsp 65 gene fragment with Xho I to produce restriction fragment(s), and analyzing the restriction fragment(s) according to an RFLP analysis method to differentiate TB complex (*Mycobacterium tuberculosis* *Mycobacterium tuberculosis* complex) and MOTT (*Mycobacteria* *Mycobacteria* other than *Mycobacterium tuberculosis* *Mycobacterium tuberculosis*).

11. (Original) The method of claim 10, wherein the restriction fragments are 391-bp, 150-bp, and 103-bp fragments to identify the TB complex.

12. (Original) The method of claim 10, wherein the 644-bp hsp 65 gene fragment is not cleaved by a restriction enzyme to identify fast-growing mycobacteria of MOTT.

13. (Original) The method of claim 10, wherein the restriction fragments are 391-bp, 169-bp, and 48-bp to identify a mycobacterial species selected from the group consisting of *M. avium*, *M. intracellulare*, *M. celatum*, *M. shimoidi*, and *M. szulgai*.

14. (Original) The method of claim 10, wherein the restriction fragments are 391-bp and 253-bp to identify a mycobacterial species selected from the group consisting of *M. gastri*, *M. genavense*, *M. gordonae*, *M. haemophilum*, *M. kansasii*, *M. malmoense*, *M. marinum*, *M. scrofulaceum*, *M. simiae*, and *M. ulcerans*.

15. (Original) A kit for the differentiation or diagnosis of TB complex and MOTT comprising a pair of primers of claim 1 and Xho I, wherein the mycobacterial species is differentiated or diagnosed based on the size of restriction fragment(s) which is obtained by amplifying an hsp 65 gene fragment of mycobacterial species in a sample with the primers to produce an amplified fragment and analyzing the amplified fragment according to an RFLP analysis method.

16. (Currently Amended) A method for the identification of a mycobacterial species comprising the steps of:

(1) amplifying an hsp 65 gene fragment of a mycobacterial species of interest with primers for specifically amplifying an hsp65 gene of mycobacteria; and

(2) hybridizing the amplified hsp65 gene fragment with a probe set comprising at least a probe selected from the group of polynucleotides consisting of ~~polynucleotides as shown in~~ SEQ ID NO: 1 to SEQ ID NO: 54 and polynucleotides complementary thereto.